PubMed Health. A service of the National Library of Medicine, National Institutes of Health.

A.D.A.M. Medical Encyclopedia. Atlanta (GA): A.D.A.M.; 2011.

Obesity

Morbid obesity; Fat - obese

Last reviewed: May 12, 2012.

Obesity means having too much body fat. It is not the same as being overweight, which means weighing too much. A person may be overweight from extra muscle, bone, or water, as well as from having too much fat.

Both terms mean that a person's weight is higher than what is thought to be healthy for his or her height.

Causes, incidence, and risk factors

Taking in more calories than you burn can lead to obesity because the body stores unused calories as fat. Obesity can be caused by:

- · Eating more food than your body can use
- · Drinking too much alcohol
- · Not getting enough exercise

Many obese people who lose large amounts of weight and gain it back think it is their fault. They blame themselves for not having the willpower to keep the weight off. Many people regain more weight than they lost.

Today, we know that biology is a big reason why some people cannot keep the weight off. Some people who live in the same place and eat the same foods become obese, while others do not. Our bodies have a complex system to help keep our weight at a healthy level. In some people, this system does not work normally.

Other factors that affect weight include:

- The way we eat when we are children can affect the way we eat as adults. The way we eat over many
 years becomes a habit. It affects what we eat, when we eat, and how much we eat.
- · We are surrounded by things that make it easy to overeat and hard to stay active:
 - o Many people do not have time to plan and make healthy meals.
 - More people today work desk jobs compared to more active jobs in the past.
 - o People with less free time have less time to exercise.

The term "eating disorder" means a group of medical conditions that have an unhealthy focus on eating, dieting, losing or gaining weight, and body image. A person may be obese, follow an unhealthy diet, and have an eating disorder all at the same time.

Sometimes, medical problems or treatments cause weight gain, including:

- Underactive thyroid gland (hypothyroidism)
- · Medicines such as birth control pills, antidepressants, and antipsychotics

Other things that can cause weight gain are:

- Quitting smoking. Most people who quit smoking gain 4 10 pounds in the first 6 months after quitting.
 Some people gain as much as 25 30 pounds.
- Stress, anxiety, feeling sad, or not sleeping well
- For women:
 - o Menopause -- women may gain 12-15 pounds during menopause
 - o Not losing the weight they gained during pregnancy

Signs and tests

The health care provider will perform a physical exam and ask questions about your medical history, eating

1 of 3 12/30/2012 10:59 PM

Obesity - PubMed Health

habits, and exercise routine.

The two most common ways to assess your weight and measure health risks related to your weight are:

- Body mass index (BMI)
- Waist circumference (your waist measurement in inches)

BMI is calculated using height and weight. You and your health care provider can use your BMI to estimate how much body fat you have.

Your waist measurement is another way to estimate how much body fat you have. Extra weight around your middle or stomach area increases your risk for type 2 diabetes, heart disease, and stroke. People with "apple-shaped" bodies (meaning their waist is bigger than their hips) also have an increased risk for these diseases.

Skin fold measurements may be taken to check your body fat percentage.

Blood tests may be done to look for thyroid or hormone problems that could lead to weight gain. However, these tests are usually normal.

Treatment

CHANGING YOUR LIFESTYLE

An active lifestyle and plenty of exercise, along with healthy eating, is the safest way to lose weight. Even modest weight loss can improve your health. You will need a lot of support from family and friends.

When dieting, your main goal should be to learn new, healthy ways of eating and make them a part of your daily routine.

Many people find it hard to change their eating habits and behaviors. You may have practiced some habits for so long that you may not even know they are unhealthy, or you do them without thinking. You need to be motivated to make lifestyle changes. Make the behavior change part of your life over the long term. Know that it takes time to make and keep a change in your lifestyle.

Work with your health care provider and dietitian to set realistic, safe daily calorie counts that help you lose weight while staying healthy. Remember that if you drop pounds slowly and steadily, you are more likely to keep them off. Your dietitian can teach you about:

- · Healthy food choices
- · Healthy snacks
- · How to read nutrition labels
- · New ways to prepare food
- Portion sizes
- Sweetened drinks

Extreme diets (fewer than 1,100 calories per day) are not thought to be safe or to work very well. These types of diets often do not contain enough vitamins and minerals. Most people who lose weight this way return to overeating and become obese again.

Learn new ways to manage stress, rather than snacking. Examples may be meditation, yoga, or exercise. If you are depressed or stressed a lot, talk to your health care provider.

MEDICATIONS AND HERBAL REMEDIES

You may see ads for supplements and herbal remedies that claim they will help you lose weight. Many of these claims are not true. Some of these supplements can have serious side effects. Talk to your health care provider before using them.

You can discuss weight loss drugs with your doctor. Most people lose between 5 and 10 pounds by taking these drugs. Most people also regain the weight when they stop taking the medicine, unless they have made lasting lifestyle changes.

SURGERY

Bariatric surgery can reduce the risk of disease in people with severe obesity. These risks include:

2 of 3 12/30/2012 10:59 PM

- Arthritis
- Diabetes
- · Heart disease
- High blood pressure
- · Sleep apnea
- · Some cancers
- Stroke

Surgery may help people who have been very obese for 5 years or more and have not lost weight from other treatments, such as diet, exercise, or medicine.

Surgery alone is not the answer for weight loss. It can train you to eat less, but you still have to do much of the work. You must be committed to <u>diet and exercise after surgery</u>. Talk to your doctor to learn if this is a good option for you.

Weight-loss surgeries include:

- · Laparoscopic gastric banding
- Gastric bypass surgery
- Sleeve gastrectomy (less common)

Support Groups

Many people find it easier to follow a diet and exercise program if they join a group of people with similar problems.

Complications

Obesity is a major health threat. The extra weight puts added stress on every part of your body and creates many risks to your health.

References

- 1. Dale KS, McAuley KA, Taylor RW, Williams SM, Farmer VL, Hansen P, et al. Determining optimal approaches for weight maintenance: a randomized controlled trial. *CMAJ*. 2009;180:E39-E46.
- 2. Jensen MD. Obesity. In: Goldman L, Schafer AI, eds. *Cecil Medicine*. 24th ed. Philadelphia, Pa: Saunders Elsevier; 2011:chap 227.
- 3. Schauer PR, Kashyap SR, Wolski K, et al. Bariatric surgery versus intensive medical therapy in obese patietns with diabetes. *N Engl J Med*. 2012:Epub March 26.
- 4. Seagle HM, Strain GW, Makris A, Reeves RS; American Dietetic Association. Position of the American Dietetic Association: weight management. *J Am Diet Assoc*. 2009;109:330-346.
- United States Department of Agriculture. Dietary Guidelines for Americans, 2010. Accessed April 21, 2012.
- 6. Wadden TA, Volger S, Sarwer DB, et al. A two year randomized trial of obesity treatment in primary care practice. *N Engl J Med*. 2011;365:1969-1979.

Review Date: 5/12/2012.

Reviewed by: David C. Dugdale, III, MD, Professor of Medicine, Division of General Medicine, Department of Medicine, University of Washington School of Medicine; and Linda J. Vorvick, MD, Medical Director and Director of Didactic Curriculum, MEDEX Northwest Division of Physician Assistant Studies, Department of Family Medicine, UW Medicine, School of Medicine, University of Washington. Also reviewed by David Zieve, MD, MHA, Medical Director, A.D.A.M. Health Solutions, Ebix, Inc.



A.D.A.M., Disclaimer

Copyright © 2012, A.D.A.M., Inc.

3 of 3 12/30/2012 10:59 PM